

# Estimating Econometric Models through Matrix Equations

Federico Poloni<sup>1</sup>  
Giacomo Sbrana<sup>2</sup>

<sup>1</sup>U Pisa, Dept of Computer Science

<sup>2</sup>Rouen Business School, France

No Free Lunch Seminar  
SNS, Pisa, February 2013

# VARMA(1,1) models

## VARMA(1,1)

$$x_t - \Phi x_{t-1} = u_t - \Theta u_{t-1}$$

$x_t$  = observed variable  $\in \mathbb{R}^d$

$u_t$  = white noise (enough to assume **uncorrelated**)  $\in \mathbb{R}^d$

$\Phi, \Theta \in \mathbb{R}^{d \times d}$ ,  $\rho(\Phi) < 1$ ,  $\rho(\Theta) < 1$

Many known models to simulate volatility reduce to VARMA(1,1):

- GARCH(1,1)
- Multivariate stochastic volatility models

# Estimating VARMA

## Problem

Given enough observations  $(x_t)$  generated by a VARMA, determine parameters  $\Phi, \Theta$

A common choice is **QML** (quasi-maximum-likelihood):

- 1 Assume  $u_t$  Gaussian independent
- 2 Given guesses  $\hat{\Phi}, \hat{\Theta}$ , compute **likelihood**  $\ell(\hat{\Phi}, \hat{\Theta})$  of generating the given time series
- 3 Feed  $\ell(\cdot, \cdot)$  into a black-box minimization procedure (e.g., Matlab's `fminunc`)

# Problems with QML

- ① Costly: each function evaluation costs  $O(nd^3)$ , with  $n =$  length of time series. Hundreds or thousands required
- ② Black-box: difficult to implement and tweak, and understand what's going on.
- ③ No convergence guarantees, non-convex optimization problem in many variables
- ④ Hey doc, what if our  $u_t$  isn't Gaussian independent?

## Our attempt

Moments estimator: determine  $\Phi, \Theta$  as a function of the **autocovariances**

$$M_k = \mathbb{E}_t \left[ x_t x_{t+k}^T \right]$$

( $\mathbb{E}_t [\cdot]$  := stationary limit mean)

The  $M_k$  contain cov's among all variables of the time series, e.g., ( $d = 3$ )

$$x_t = \begin{bmatrix} a_t \\ b_t \\ c_t \end{bmatrix} \Rightarrow M_k = \begin{bmatrix} \mathbb{E}_t [a_t a_{t+k}] & \mathbb{E}_t [a_t b_{t+k}] & \mathbb{E}_t [a_t c_{t+k}] \\ \mathbb{E}_t [b_t a_{t+k}] & \mathbb{E}_t [b_t b_{t+k}] & \mathbb{E}_t [b_t c_{t+k}] \\ \mathbb{E}_t [c_t a_{t+k}] & \mathbb{E}_t [c_t b_{t+k}] & \mathbb{E}_t [c_t c_{t+k}] \end{bmatrix}$$

# Moment estimator

We will show  $(\Phi, \Theta) = f(M_0, M_1, M_2)$

## GMM estimator

- 1 Compute sample  $\hat{M}_k = \frac{1}{n} \sum x_t x_{t+k}^T$
  - 2 Get  $(\hat{\Phi}, \hat{\Theta}) = f(\hat{M}_0, \hat{M}_1, \hat{M}_2)$
- Very fast: working only with  $d \times d$  matrices, no dependence on  $n$  (after computing moments)
  - Asymptotically consistent and normal, under suitable conditions
  - In simulated experiments, not as accurate as QML, but good as initial value

Already known for **univariate GARCH**; generalization requires some linear algebra machinery

# Yule-Walker results

The parameter  $\Phi$  is easy to obtain:

## Theorem

$$\Phi = M_{k+1}M_k^{-1} \text{ for each } k \geq 1$$

In particular  $\Phi = M_2M_1^{-1}$

Proof:

$$\underbrace{x_{t-k-1}(x_t - \Phi x_{t-1})^T}_{=M_{k+1}-\Phi M_k} = \underbrace{x_{t-k-1}(u_t - \Theta u_{t-1})^T}_{=0 \text{ if } k \geq 1}$$

## Estimating $\Theta$

Let  $r_t = x_t - \Phi x_{t-1} = u_t - \Theta u_{t-1}$ ,  $Y := \mathbb{E} [u_t u_t^T]$

$$\Gamma_0 := \mathbb{E}_t [r_t r_t^T] = M_0 - \Phi M_1^T - M_1 \Phi^T + \Phi M_0 \Phi^T = Y + \Theta Y \Theta^T$$

$$\Gamma_1 := \mathbb{E}_t [r_t r_{t+1}^T] = M_1 - \Phi M_0 = -\Theta Y$$

Blue expressions allow us to compute  $\Gamma_0$ ,  $\Gamma_1$ .

Use them + red expressions to decouple equations for  $Y$ ,  $X = \Theta^T$

$$\begin{aligned}\Gamma_0 &= Y + \Gamma_1 Y^{-1} \Gamma_1^T \\ \Gamma_1^T + \Gamma_0 X + \Gamma_1 X^2 &= 0\end{aligned}$$



## Existence and unicity

$$\begin{aligned} \Gamma_0 &= Y + \Gamma_1 Y^{-1} \Gamma_1^T \\ \Gamma_1^T + \Gamma_0 X + \Gamma_1 X^2 &= 0 \end{aligned}$$

Studied by matrix equation people (like me)

### Existence and unicity

- Solution exists if  $Q(\lambda) := \Gamma_1^T \lambda^{-1} + \Gamma_0 + \Gamma_1 \lambda$  is such that  $Q(\lambda) > 0$  for each  $\lambda$  on unit circle
- Solution unique if we ask  $Y > 0$ ,  $\rho(X) < 1$  (as was assumed)

Of course, if the model is well-posed, there must be a solution. . .

# How to solve matrix equations

Let us focus on  $\Gamma_1^T + \Gamma_0 X + \Gamma_1 X^2 = 0$

Solution resembles a lot linear recurrence theory

Generalized **eigenvalues/Vectors** of the problem:

$$(\lambda, v) \quad \text{s.t.} \quad (\Gamma_1^T + \Gamma_0 \lambda + \Gamma_1 \lambda^2)v = 0$$

(there are  $2d$  of them!)

## Theorem

Solutions can be eigendecomposed as  $X = VDV^{-1}$

$V$  contains  $d$  of the  $2d$  eigenvectors of the problem,  $D$  diagonal with eigenvalues

# Generalized eigenvalues

## Companion matrix

Eigenvalues/eigenvectors are in 1:1 correspondence with those of the **linearization matrix**

$$\begin{bmatrix} 0 & I_d \\ -\Gamma_1^{-1}\Gamma_1^T & -\Gamma_1^{-1}\Gamma_0 \end{bmatrix}$$

Matrix version of the “companion matrix” for polynomials

## Palindromic matrix polynomials

Due to the structure in  $\Gamma_1^T + \Gamma_0\lambda + \Gamma_1\lambda^2$ ,  $\Gamma_0 = \Gamma_0^T$ , **eigenvalues come in pairs**  $(\lambda, \lambda^{-1})$

Good for us — we needed  $d$  of them with  $|\lambda| < 1$ !

## To sum up

- 1 Get sample moments  $M_0, M_1, M_2$
- 2 Get  $\Phi = M_2 M_1^{-1}$
- 3 Compute  $\Gamma_0, \Gamma_1$
- 4 Get eigenvalues/vectors of  $\begin{bmatrix} 0 & I_d \\ -\Gamma_1^{-1} \Gamma_1^T & -\Gamma_1^{-1} \Gamma_0 \end{bmatrix}$
- 5 Take those with  $\lambda$  inside the unit circle
- 6 Assemble  $\Theta^T = X = V D V^{-1}$

## What can go wrong


Sometimes, **no stationary/invertible model** with autocovariances  $\hat{M}_i$

### Existence and unicity

- Solution  $X, Y$  exists if  $Q(\lambda) := \Gamma_1^T \lambda^{-1} + \Gamma_0 + \Gamma_1 \lambda$  is such that  $Q(\lambda) > 0$  for each  $\lambda$  on unit circle

Must hold with **exact**  $\Gamma_i$ , but **sample**  $\hat{\Gamma}_i$  might give inconsistent data

$\Rightarrow$  Perturb  $M_0, M_1, M_2$  to ensure solvability!



## Some results




Figure: Diagonal GARCH,  $d = 2$ ,  $\rho(\Theta) = 0.6$ ,  $n = 1000$

## Some results




Figure: Diagonal GARCH,  $d = 2$ ,  $\rho(\Theta) = 0.6$ ,  $n = 1000$

## Some results




Figure: Diagonal GARCH,  $d = 2$ ,  $\rho(\Theta) = 0.7$ ,  $n = 1000$



## Some results




Figure: Diagonal GARCH,  $d = 3$ ,  $\rho(\Theta) = 0.6$ ,  $n = 1000$

## Some results




Figure: Diagonal GARCH,  $d = 3$ ,  $\rho(\theta) = 0.6$ ,  $n = 1000$ , larger off-diagonal elements

## Some results




Figure: Diagonal GARCH,  $d = 3$ ,  $\rho(\Theta) = 0.6$ ,  $n = 500$

## Some results




Figure: Diagonal GARCH,  $d = 2$ ,  $\rho(\Theta) = 0.7$ ,  $n = 500$

## Some results




Figure: Diagonal GARCH,  $d = 2$ ,  $\rho(\theta) = 0.6$ ,  $n = 5000$

## Some results




Figure: Diagonal GARCH,  $d = 2$ ,  $\rho(\Theta) = 0.7$ ,  $n = 5000$

## Possible improvements

- Improve solvability enforcement (work on  $\Theta$  and  $\Phi$  at the same time)
- Combine with an iterative ML-like optimization  
e.g., GLS (generalized least squares) for GARCH?
- More intensive testing & applications

## Possible improvements

- Improve solvability enforcement (work on  $\Theta$  and  $\Phi$  at the same time)
- Combine with an iterative ML-like optimization  
e.g., GLS (generalized least squares) for GARCH?
- More intensive testing & applications

Thanks for your attention!